arXiv:0911.1977v2 [gr-qc] 11 Nov 2009

Solving the Einstein field equations

A.V.Bratchikov
Kuban State Technological University,
2 Moskovskaya Street, Krasnodar, 350072, Russia

Abstract

A solution of the Einstein vacuum field equations is constructed within the contex of perturbation theory. The solution possesses a graphical representation in terms of diagrams.

In a recent article [1] for a wide class of nonlinear equations a perturbative solution was constructed.This class includes equations of motion of field theories. Solutions of the pure Yang-Mills field equations and equations of motions in spinor electrodynamics [2] were obtained. The solutions can be described by diagrams. The aim of the present paper is to construct a solution of the Einstein field equations in the vacuum.

Let $x = (x^0, x^1, x^2, x^3)$ be space-time coordinates, and $g(x) = g_{\mu\nu}(x)dx^\mu dx^\nu$ a space-time metric. The vacuum Einstein equations read

$$G_{\mu\nu} = 0, \tag{1}$$

where

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R,$$

$R_{\mu\nu}$ is the Ricci tensor

$$R_{\mu\nu} = \Gamma^\beta_{\mu\nu,\beta} - \Gamma^\beta_{\mu\beta,\nu} + \Gamma^\beta_{\mu\nu}\Gamma^\alpha_{\beta\alpha} - \Gamma^\alpha_{\mu\beta}\Gamma^\beta_{\nu\alpha},$$

$R = R^\mu_\mu$, and $\Gamma^\gamma_{\alpha\beta}$ is the Cristofel symbol

$$\Gamma^\gamma_{\alpha\beta} = \frac{1}{2} g^{\gamma\rho}(\partial_\beta g_{\rho\alpha} + \partial_\alpha g_{\rho\beta} - \partial_\rho g_{\alpha\beta}).$$

For $\eta = \eta_{\mu\nu} dx^\mu dx^\nu, \eta_{\mu\nu} = diag(1, -1, -1, -1)$, we can expand equations (1) in powers of $h = g - \eta$

$$\sum_{n=1}^{\infty} G^n_{\mu\nu}(h) = 0. \tag{2}$$

Here

$$G^n_{\mu\nu}(h) = \frac{1}{n!} \frac{d^n G_{\mu\nu}(\eta + \xi h)}{d^n \xi}\bigg|_{\xi=0}.$$

We impose a gauge condition

$$\eta^{\alpha\beta} \partial_\alpha \varphi_{\beta\nu} = 0, \tag{3}$$

where

$$\varphi_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu} h^\alpha_\alpha, \qquad h^\alpha_\alpha = \eta^{\alpha\beta} h_{\alpha\beta}. \tag{4}$$

From equation (4) it follows

$$h_{\mu\nu} = \varphi_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu} \varphi^\alpha_\alpha, \qquad \varphi^\alpha_\alpha = \eta^{\alpha\beta} \varphi_{\alpha\beta}. \tag{5}$$

Let $\tilde{G}^n(\varphi)$ be defined by $\tilde{G}^n(\varphi) = G^n(h)_{\mu\nu} dx^\mu dx^\nu$, $\varphi = \varphi_{\mu\nu} dx^\mu dx^\nu$. Then equation (2) takes the form [3]

$$\Box \varphi - 2 \sum_{n=2}^{\infty} \tilde{G}^n(\varphi) = 0, \tag{6}$$

where $\Box = \eta^{\alpha\beta} \partial_\alpha \partial_\beta$.

The general solution $\varphi_{0\mu\nu}$ of the free equation

$$\Box \varphi_{\mu\nu} = 0$$

is given by [4]

$$\varphi_{0\mu\nu} = \frac{1}{4\pi} M(v_{\mu\nu}) + \frac{1}{4\pi} \frac{\partial}{\partial t}(t M(u_{\mu\nu})),$$

where

$$M(w_{\mu\nu}) = \int_S w_{\mu\nu}(x^1 + t\xi^1, x^2 + t\xi^2, x^3 + t\xi^3) d\sigma_\xi,$$

ξ^1, ξ^2, ξ^3 are coordinates on the unit sphere S, σ_ξ is the area element on S, $t = x^0$,

$$\varphi_{0\mu\nu}(0, x^1, x^2, x^3) = u_{\mu\nu}(x^1, x^2, x^3),$$

$$\left. \frac{\partial \varphi_{0\mu\nu}(t, x^1, x^2, x^3)}{\partial t} \right|_{t=0} = v_{\mu\nu}(x^1, x^2, x^3),$$

and $u_{\mu\nu}, v_{\mu\nu}$ are some functions. Equation (3) implies

$$v_{0\nu} - \sum_{i=1}^{3} \partial_i u_{i\nu} = 0.$$

A specific solution $\mathcal{E}(f_{\mu\nu})$ of the inhomogeneous equation

$$\Box \varphi_{\mu\nu} = f_{\mu\nu}$$

reads

$$\mathcal{E}(f_{\mu\nu}) = \frac{1}{4\pi} \int_0^t \tau d\tau \int_S f_{\mu\nu}(t - \tau, x^1 + \tau\xi^1, x^2 + \tau\xi^2, x^3 + \tau\xi^3) d\sigma_\xi.$$

Then equation (6) can be written in the form

$$\varphi = \kappa\varphi_0 + \sum_{n=2}^{\infty} p_n(\varphi), \tag{7}$$

where $\varphi_0 = \varphi_{0\mu\nu} dx^\mu dx^\nu$, and $p_n = 2\mathcal{E}(\tilde{G}^n_{\mu\nu}) dx^\mu dx^\nu$. Here we introduced an auxiliary parameter κ. Eventually it is set to be 1.

Let V be the space of symmetric 2-forms. We shall need the polylinear symmetric functions

$$\langle \ldots \rangle_n : V^n \to V, \quad n = 2, 3, \ldots,$$

defined for $\gamma_1, \ldots, \gamma_n \in V$ by

$$\langle \gamma_1, \ldots, \gamma_n \rangle_n = \frac{\partial}{\partial a_1} \ldots \frac{\partial}{\partial a_n} p_n(a_1\gamma_1 + \ldots + a_n\gamma_n). \tag{8}$$

One can check that

$$\langle \gamma, \ldots, \gamma \rangle_n = n! p_n(\gamma).$$

Then equation (7) becomes

$$\varphi = \kappa\varphi_0 + \sum_{n=2}^{\infty} \frac{1}{n!} \langle \varphi, \dots, \varphi \rangle_n. \tag{9}$$

For $m \geq 2, 1 \leq i_1 < \dots < i_n \leq m$ let

$$P^m_{i_1 \dots i_n} : V^m \to V^{m-n+1}$$

be defined by

$$P^m_{i_1 \dots i_n}(\gamma_1, \dots, \gamma_m) = (\langle \gamma_{i_1}, \dots, \gamma_{i_n} \rangle_n, \gamma_1, \dots, \widehat{\gamma}_{i_1}, \dots, \widehat{\gamma}_{i_n}, \dots, \gamma_m),$$

where $\widehat{\gamma}$ means that γ is omitted. If $\phi \in V$ is given by

$$\phi = P^{n_s}_{I_s} \dots P^{m-n_1+1}_{I_2} P^m_{I_1}(\gamma_1, \dots, \gamma_m) \tag{10}$$

for some $I_1 = (i^1_1 \dots i^1_{n_1}), I_2 = (i^2_1 \dots i^2_{n_2}), \dots, I_s = (i^s_1 \dots i^s_{n_s})$, we say that ϕ is a descendant of $(\gamma_1, \dots, \gamma_m)$.

Each descendant can be represented by a diagram. In this diagram an element of V is represented by the line segment —— . A product

$$(\gamma_1, \dots, \gamma_n) \to \langle \gamma_1, \dots, \gamma_n \rangle_n$$

is represented by the vertex joining the line segments for $\gamma_1, \dots, \gamma_n$, $\langle \gamma_1, \dots, \gamma_n \rangle_n$. The general rule for graphical representation of $P^m_I(\gamma_1, \dots, \gamma_m)$ should be clear from Figure 1.Here we show the diagram for

$$P^m_{ijk}(\gamma_1, \dots, \gamma_m) = (\langle \gamma_i, \gamma_j, \gamma_k \rangle_3, \gamma_1, \dots, \widehat{\gamma}_i, \dots, \widehat{\gamma}_j, \dots, \widehat{\gamma}_k, \dots, \gamma_m).$$

The points labeled by $1, \dots, m$ represent the ends of the lines for $\gamma_1, \dots, \gamma_m$. Using the representation for $P^m_{i_1 \dots i_n}(\gamma_1, \dots, \gamma_m)$ one can consecutively draw the diagrams for $P^{n_1}_{I_1}(\gamma_1, \dots, \gamma_m), P^{n_2}_{I_2} P^{n_1}_{I_1}(\gamma_1, \dots, \gamma_m), \dots, \phi$ (10).

Let us introduce a family of functions

$$\langle \dots \rangle : V^m \to V, \qquad m = 2, 3, \dots,$$

such that for $\gamma_1, \dots, \gamma_m \in V$ $\langle \gamma_1, \dots, \gamma_m \rangle$ is defined as the sum of all the descendants of its arguments. For example, for $m = 2$ and $m = 3$ we have

$$\langle \gamma_1, \gamma_2 \rangle = \langle \gamma_1, \gamma_2 \rangle_2,$$



Figure 1. Diagram for $P^m_{ijk}(\gamma_1,\ldots,\gamma_m)$.

$$\langle\gamma_1,\gamma_2,\gamma_3\rangle=\langle\langle\gamma_1,\gamma_2\rangle_2,\gamma_3\rangle_2+\langle\langle\gamma_1,\gamma_3\rangle_2,\gamma_2\rangle_2+\langle\langle\gamma_2,\gamma_3\rangle_2,\gamma_1\rangle_2+\langle\gamma_1,\gamma_2,\gamma_3\rangle_3.$$

A solution of equation (9) is given by [1]

$$\varphi=\langle e^{\kappa\varphi_0}\rangle,$$

where

$$\langle e^{\kappa\varphi_0}\rangle=\sum_{n=0}^{\infty}\frac{\kappa^n}{n!}\langle\varphi_0^n\rangle,$$

$\langle\varphi_0^n\rangle$ is defined as 0 if $n=0$, and otherwise

$$\langle\varphi_0^n\rangle=\langle\underbrace{\varphi_0,\ldots,\varphi_0}_{\text{n times}}\rangle,\qquad\langle\varphi_0\rangle=\varphi_0.$$

For example, the $O(\kappa^3)$ contribution in φ is given by

$$\frac{\kappa^3}{6}\langle\varphi_0,\varphi_0,\varphi_0\rangle=\kappa^3\left(\frac{1}{2}\langle\langle\varphi_0,\varphi_0\rangle_2,\varphi_0\rangle_2+\frac{1}{6}\langle\varphi_0,\varphi_0,\varphi_0\rangle_3\right).$$

The diagram for $\langle\langle\varphi_0,\varphi_0\rangle_2,\varphi_0\rangle_2$ is depicted in Figure 2.


φ_0
φ_0
φ_0
$\langle\langle\varphi_0, \varphi_0\rangle_2, \varphi_0\rangle_2$

Figure 2. Diagram for $\langle\langle\varphi_0, \varphi_0\rangle_2, \varphi_0\rangle_2$.

The tensor $h_{\mu\nu}$ can be found by using (5).

References

[1] A.V.Bratchikov, Perturbation theory for nonlinear equations, arXiv 0910.3924 [hep-th].

[2] A.V.Bratchikov, Solving field equations in spinor electrodynamics, arXiv 0910.0621 [hep-th].

[3] L.D.Landau and D.V.Lifshits, The classical theory of fields, Nauka, Moscow, 1973.

[4] R.Courant and D.Hilbert, Methods of mathematical physics, Wiley, 1989.